



Adult and Pediatric Urology

BLOOD THINNERS

- BLOOD THINNING MEDICATIONS SHOULD BE DISCONTINUED **5 DAYS** BEFORE HAVING SURGERY OR AS DIRECTED BY YOUR SURGEON.
- BE SURE TO TALK WITH YOUR DOCTOR BEFORE RESUMING THESE MEDICATIONS OR IF YOU HAVE ANY QUESTIONS REGARDING YOUR CURRENT MEDICATIONS.

ADVIL
ALEVE
ALKA SELTZER
ARIXTRA
ANAPROX
ASPIRIN
BEXTRA
CELEBREX
COUMADIN
FISH OIL
IBUPROFEN
MOTRIN
NAPROSYN
PLAVIX
TICLID
VIOXX
VITAMIN E
VOLTAREN

Karl R. Westenfelder, M.D., Garrick R. Simmons, M.D. Kevin M. Kronner, M.D., Christopher G Wicher, M.D.
North Campus - Garden City Medical Arts Building 601 W. Spruce Suite G, Missoula, Montana 59802
(406) 728-3366 (800) 370-3364 Fax (406) 329-2896
South Campus – 2831 Fort Missoula Road, Physician Center II, STE 105, Missoula, MT 59804
(406) 830-3107 Fax (406) 830-3109